



THE UNIVERSITY of EDINBURGH

## Edinburgh Research Explorer

### First community-wide, comparative cross-linking mass spectrometry study

**Citation for published version:**

Iacobucci, C, Piotrowski, C, Aebersold, R, Amaral, BC, Andrews, P, Bernfur, K, Borchers, C, Brodie, NI, Bruce, JE, Cao, Y, Chaignepain, S, Chavez, JD, Claverol, S, Cox, J, Davis, T, Degliesposti, G, Dong, MQ, Edinger, N, Emanuelsson, C, Gay, M, Götze, M, Gomes-Neto, F, Gozzo, FC, Gutierrez, C, Haupt, C, Heck, AJR, Herzog, F, Huang, L, Hoopmann, MR, Kalisman, N, Klykov, O, Kukaka, Z, Liu, F, Maccoss, MJ, Mechtler, K, Mesika, R, Moritz, RL, Nagaraj, N, Nesati, V, Neves-Ferreira, AGC, Ninnis, R, Novák, P, O'Reilly, FJ, Pelzing, M, Petrotchenko, E, Piersimoni, L, Plasencia, M, Pukala, T, Rand, KD, Rappsilber, J, Reichmann, D, Sailer, C, Sarnowski, CP, Scheltema, RA, Schmidt, C, Schriemer, DC, Shi, Y, Skehel, JM, Slavin, M, Sobott, F, Solis-Mezarino, V, Stephanowitz, H, Stengel, F, Stieger, CE, Trabjerg, E, Trnka, M, Vilaseca, M, Viner, R, Xiang, Y, Yilmaz, S, Zelter, A, Ziemianowicz, D, Leitner, A & Sinz, A 2019, 'First community-wide, comparative cross-linking mass spectrometry study', *Analytical Chemistry*, vol. 91, no. 11, pp. 6953-6961. <https://doi.org/10.1021/acs.analchem.9b00658>

**Digital Object Identifier (DOI):**

[10.1021/acs.analchem.9b00658](https://doi.org/10.1021/acs.analchem.9b00658)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Analytical Chemistry

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# The First Community-Wide, Comparative Cross-linking Mass Spectrometry Study

Claudio Iacobucci<sup>1#</sup>, Christine Piotrowski<sup>1#</sup>, Ruedi Aebersold<sup>2,3</sup>, Bruno C. Amaral<sup>4</sup>, Philip Andrews<sup>5</sup>, Katja Bernfur<sup>17</sup>, Christoph Borchers<sup>6,7,8,9</sup>, Nicolas I. Brodie<sup>6</sup>, James E. Bruce<sup>10</sup>, Yong Cao<sup>15</sup>, Stéphane Chaignepain<sup>11</sup>, Juan D. Chavez<sup>10</sup>, Stéphane Claverol<sup>12</sup>, Jürgen Cox<sup>13</sup>, Trisha Davis<sup>41</sup>, Gianluca Degliesposti<sup>14</sup>, Meng-Qiu Dong<sup>15</sup>, Nufar Edinger<sup>16</sup>, Cecilia Emanuelsson<sup>17</sup>, Marina Gay<sup>18</sup>, Michael Götze<sup>19</sup>, Francisco Gomes-Neto<sup>44</sup>, Fabio C. Gozzo<sup>4</sup>, Craig Gutierrez<sup>20</sup>, Caroline Haupt<sup>21</sup>, Albert J. R. Heck<sup>22</sup>, Franz Herzog<sup>23</sup>, Lan Huang<sup>20</sup>, Michael R. Hoopmann<sup>24</sup>, Nir Kalisman<sup>16</sup>, Oleg Klykov<sup>22</sup>, Zdeněk Kukačka<sup>25</sup>, Fan Liu<sup>26</sup>, Michael J. MacCoss<sup>27</sup>, Karl Mechtler<sup>28</sup>, Ravit Mesika<sup>16</sup>, Robert L. Moritz<sup>24</sup>, Nagarjuna Nagaraj<sup>29</sup>, Victor Nesati<sup>30</sup>, Ana G. C. Neves-Ferreira<sup>44</sup>, Robert Ninnis<sup>30</sup>, Petr Novák<sup>25</sup>, Francis J O'Reilly<sup>31</sup>, Matthias Pelzing<sup>30</sup>, Evgeniy Petrotchenko<sup>6</sup>, Lolita Piersimoni<sup>5</sup>, Manolo Plasencia<sup>5</sup>, Tara Pukala<sup>32</sup>, Kasper D. Rand<sup>33</sup>, Juri Rappsilber<sup>31,34</sup>, Dana Reichmann<sup>16</sup>, Carolin Sailer<sup>35</sup>, Chris P. Sarnowski<sup>2,36</sup>, Richard A. Scheltema<sup>22</sup>, Carla Schmidt<sup>21</sup>, David C. Schriemer<sup>37</sup>, Yi Shi<sup>40</sup>, J. Mark Skehel<sup>14</sup>, Moriya Slavin<sup>16</sup>, Frank Sobott<sup>42,43</sup>, Victor Solis-Mezarino<sup>23</sup>, Heike Stephanowitz<sup>26</sup>, Florian Stengel<sup>35</sup>, Christian E. Stieger<sup>28</sup>, Esben Trabjerg<sup>33</sup>, Michael Trnka<sup>38</sup>, Marta Vilaseca<sup>18</sup>, Rosa Viner<sup>39</sup>, Yufei Xiang<sup>40</sup>, Sule Yilmaz<sup>13</sup>, Alex Zelter<sup>41</sup>, Daniel Ziemianowicz<sup>37</sup>, Alexander Leitner<sup>2\*</sup>, Andrea Sinz<sup>1\*</sup>

<sup>1</sup> Department of Pharmaceutical Chemistry and Bioanalytics, Institute of Pharmacy, Charles Tanford Protein Center, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Str. 3a, 06120 Halle/Saale, Germany.

<sup>2</sup> Department of Biology, Institute of Molecular Systems Biology, ETH Zurich, Otto-Stern-Weg 3, 8093 Zurich, Switzerland.

<sup>3</sup> Faculty of Science, University of Zurich, 8006 Zurich, Switzerland.

<sup>4</sup> Institute of Chemistry, University of Campinas, Campinas SP, 13083-970, Brazil.

<sup>5</sup> Departments of Biological Chemistry, Bioinformatics, and Chemistry, University of Michigan, Ann Arbor Michigan, MI 48109, USA.

<sup>6</sup> University of Victoria - Genome British Columbia Proteomics Centre, Vancouver Island Technology Park, Victoria, British Columbia V8Z 7X8, Canada.

<sup>7</sup> Department of Biochemistry and Microbiology, University of Victoria, Petch Building, Room 270d, 3800 Finnerty Road, Victoria, British Columbia V8P 5C2, Canada.

<sup>8</sup> Gerald Bronfman Department of Oncology, Jewish General Hospital, McGill University, 3755 Côte Ste-Catherine Road, Montréal, Quebec H3T 1E2, Canada.

<sup>9</sup> Proteomics Centre, Segal Cancer Centre, Lady Davis Institute, Jewish General Hospital, McGill University, 3755 Côte Ste-Catherine Road, Montréal, Quebec H3T 1E2, Canada.

<sup>10</sup> Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA.

<sup>11</sup> CBMN, UMR 5248, CNRS, Université de Bordeaux, INP Bordeaux, Pessac 33607, France.

<sup>12</sup> Centre de Génomique Fonctionnelle, Plateforme Protéome, Université de Bordeaux, Bordeaux 33000, France.

<sup>13</sup> Computational Systems Biochemistry Research Group, Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany.

<sup>14</sup> MRC Laboratory of Molecular Biology, Cambridge Biomedical Campus, Francis Crick Ave, Cambridge CB2 0QH, UK.

<sup>15</sup> National Institute of Biological Sciences, Beijing, 7 Science Park Road, ZGC Life Science Park, 102206 Beijing, China.

<sup>16</sup> Department of Biological Chemistry, The Alexander Silberman Institute of Life Sciences, Safra Campus Givat Ram, The Hebrew University of Jerusalem, Jerusalem 91904, Israel.

<sup>17</sup> Department of Biochemistry and Structural Biology, Center for Molecular Protein Science, Lund University, 221 00 Lund, Sweden.

- <sup>18</sup> Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science and Technology (BIST), Baldiri Reixac 10, 08028 Barcelona, Spain.
- <sup>19</sup> Institute for Biochemistry and Biotechnology, Charles Tanford Protein Center, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Str. 3a, 06120 Halle/Saale, Germany.
- <sup>20</sup> Department of Physiology & Biophysics, University of California, Irvine, CA 92697, USA.
- <sup>21</sup> Interdisciplinary Research Center HALOmeme, Institute for Biochemistry and Biotechnology, Charles Tanford Protein Center, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Str. 3a, 06120 Halle/Saale, Germany.
- <sup>22</sup> Biomolecular Mass Spectrometry and Proteomics, Bijvoet Center for Biomolecular Research and Utrecht Institute for Pharmaceutical Sciences, University of Utrecht and Netherlands Proteomics Centre, Padualaan 8, 3584 CH Utrecht, The Netherlands.
- <sup>23</sup> Gene Center Munich, Department of Biochemistry, Faculty of Chemistry and Pharmacy, Ludwig Maximilians University of Munich, Feodor-Lynen-Str. 25, 81377 Munich, Germany.
- <sup>24</sup> Institute for Systems Biology, 401 Terry Ave N, Seattle, WA 98109, USA.
- <sup>25</sup> Institute of Microbiology, BIOCEV, Prumyslova 595, 252 50 Vestec, Czech Republic.
- <sup>26</sup> Leibniz Institute of Molecular Pharmacology (FMP), Robert-Rössle-Str. 10, 13125 Berlin, Germany.
- <sup>27</sup> Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA.
- <sup>28</sup> Protein Chemistry Facility, Research Institute of Molecular Pathology (IMP) and Institute of Molecular Biotechnology (IMBA), Vienna Biocenter (VBC), Dr. Bohr-Gasse 3, 1030 Vienna, Austria.
- <sup>29</sup> Biochemistry Core Facility, Max-Planck-Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany;
- <sup>30</sup> Analytical Biochemistry, CSL Limited, Bio21 Institute, 30 Flemington Road, 3010 Parkville, Australia.
- <sup>31</sup> Chair of Bioanalytics, Institute of Biotechnology Technische Universität Berlin, 13355 Berlin, Germany.
- <sup>32</sup> Discipline of Chemistry, Faculty of Sciences, University of Adelaide, North Terrace, Adelaide, South Australia, 5005, Australia.
- <sup>33</sup> Department of Pharmacy, University of Copenhagen, 2100 Copenhagen, Denmark.
- <sup>34</sup> Wellcome Trust Centre for Cell Biology, School of Biological Sciences, University of Edinburgh, EH9 3BF Edinburgh, UK.
- <sup>35</sup> University of Konstanz, Department of Biology, Universitätsstr. 10, 78457 Konstanz, Germany.
- <sup>36</sup> PhD Program in Systems Biology, University of Zurich and ETH Zurich, 8092 Zurich, Switzerland.
- <sup>37</sup> Dept. of Biochemistry & Molecular Biology, Robson DNA Science Centre, University of Calgary, 3330 Hospital Dr. NW, Calgary, Alberta T2N 4N1, Canada.
- <sup>38</sup> UCSF Mass Spectrometry Facility, Genentech Hall, 600 16th Street, San Francisco, CA 94158, USA.
- <sup>39</sup> Thermo Fisher Scientific, 355 River Oaks Parkway, San Jose, CA 95134, USA.
- <sup>40</sup> Department of Cell Biology, University of Pittsburgh, School of Medicine, Pittsburgh, PA 15213, USA.
- <sup>41</sup> Department of Biochemistry, University of Washington, Seattle, WA 98195, USA.
- <sup>42</sup> Department of Chemistry, University of Antwerp, Groenenborgerlaan 171, 2020 Antwerp, Belgium.
- <sup>43</sup> The Astbury Centre for Structural Molecular Biology, and School of Molecular and Cellular Biology, University of Leeds, LS2 9JT Leeds, UK.
- <sup>44</sup> Laboratory of Toxinology, Oswaldo Cruz Institute, Fiocruz, Avenida Brasil 4.365, Manguinhos, Rio de Janeiro - RJ, CEP 21040-900 (Moorish Castle), Brazil.

---

**ABSTRACT:** The number of publications in the field of chemical cross-linking combined with mass spectrometry (XL-MS) to derive constraints for protein three-dimensional structure modeling and to probe protein-protein interactions has increased during the last years. As the technique is now becoming routine for in vitro and in vivo applications in proteomics and structural biology there is a pressing need to define protocols as well as data analysis and reporting formats. Such consensus formats should become accepted in the field and be shown to lead to reproducible results. This first, community-based harmonization study on XL-MS is based on the results of 32 groups participating worldwide. The aim of this paper is to summarize the status quo of XL-MS and to compare and evaluate existing cross-linking strategies. Our study therefore builds the framework for establishing best practice guidelines for conducting cross-linking experiments, performing data analysis, and reporting formats with the ultimate goal of assisting scientists to generate accurate and reproducible XL-MS results.

---

INTRODUCTION. Mass spectrometry (MS) is becoming increasingly popular in the field of structural biology, with great

implications for solving important biological questions. A central technique in structural MS is chemical cross-linking combined with

MS (XL-MS). Since 2000, XL-MS and computational modeling has advanced from investigating three-dimensional structures of isolated proteins to deciphering protein interaction networks<sup>1-4</sup>. In the field of integrated structure analysis, XL-MS is often used in conjunction with cryo-electron microscopy. As the chemical XL-MS approach allows the capture of transient and weak interactions, it is now becoming a routine technique for unraveling protein interaction networks in their natural cellular environment<sup>5</sup>. The knowledge obtained will significantly advance our understanding of the structure of functional complexes, the topology of cellular networks and molecular details underlying human pathologies. Briefly, the XL-MS approach relies on adding a chemical reagent to a protein solution connecting two functional groups of amino acid side chains. Cross-linker molecules consist of two reactive groups that are separated via a spacer of defined length that allow to derive distance information of a protein or a protein assembly. The cross-linked residues are usually identified after enzymatic digestion of the covalently connected protein(s) using LC/ESI-MS/MS (liquid chromatography electrospray ionization tandem mass spectrometry) and the resulting fragment ion spectra are computationally assigned to the cross-linked peptides. The distance constraints imposed by the chemical cross-linker on the protein's tertiary structure serve as a basis for subsequent computational modeling studies to derive three-dimensional structural models (Scheme S1). XL-MS can be applied to both proteins and protein complexes and in the case of protein assemblies, the distance constraints can be used to map the subunit topology. XL-MS is now increasingly being used for deriving protein-protein interaction maps, both in vitro and in vivo, where interacting proteins are covalently connected by the cross-linking reaction.<sup>6-13</sup>

The wide acceptance of XL-MS by the proteomics and structural biology communities reflects the increasing importance of cross-linking data for elucidating protein structures and protein-protein interactions. However, the growth of the user base brings about challenges of its own: Even a relatively superficial glance at the literature shows a huge diversity of cross-linkers, experimental workflows, and computational pipelines. Moreover, the information provided in scientific research articles that contain cross-linking data can range from being quite detailed to very brief.

The heterogeneity of cross-linking protocols has mainly emerged from the use of different cross-linking chemistries and different designs of the corresponding cross-linker (e.g., non-cleavable/cleavable, isotope-coded, or affinity-tagged reagents). This, in turn, necessitated individual software solutions specifically tailored to the analysis of data from the experimental workflow. The most common database search engines used in proteomics are not directly suitable for interpreting mass spectra from cross-linked peptides. Therefore, the majority of computational solutions have emerged from laboratories that pioneered the application of XL-MS and created tools specifically tailored for the analysis of cross-linked peptides. Together with a current lack of formal or even informal reporting standards, the present state of XL-MS may confuse researchers that are interested in interpreting results from XL-MS studies or in adopting the technology. Currently, it is not clear, which strategies are most suitable in general or for a particular application, which makes it challenging to objectively compare results obtained by different groups.

Certainly, the challenges summarized above resemble those of other disciplines. In particular, scientists active in "conventional" proteomics research have tried to address the very same issues over the last decade. Inter-laboratory and software comparison studies have been performed for different experimental strategies, including data-dependent acquisition<sup>14</sup>, selected reaction monitoring<sup>15-18</sup>, and most recently, data-independent acquisition<sup>19,20</sup>. In addition, regular comparative studies have been organized by the Association of Biomolecular Resource Facilities (ABRF; <https://abrf.org/research-group/proteomics-research-group-prg> and <https://abrf.org/research-group/proteomics-standards-research-group-sprg>). Together, these studies revealed limitations in commonly used experimental and computational workflows, but on the other hand also provided evidence for the robustness of a particular technique when implemented in different laboratories according to standard operating procedures.

Standardized file formats and reporting guidelines for proteomics have been developed under the auspices of the Proteomics Standards Initiative (PSI) of the Human Proteome Organization (<http://www.psivdev.info>)<sup>21</sup>. For example, as far back as 2007, the first recommendations for minimum reporting standards in proteomics (Minimum Information About a Proteomics Experiment, MIAPE) have been made<sup>22</sup>, which have been followed by detailed guidelines of several proteomics journals. PSI has also formalized open-file formats, such as the mzML format for raw MS data<sup>23</sup> and the mzIdentML format for protein identifications<sup>24</sup>. Such guidelines and open data formats have also led to an increase in the deposition of proteomics data in open data repositories such as the PRoteomics IDentifications (PRIDE) archive, hosted by the European Bioinformatics Institute (<https://www.ebi.ac.uk/pride/archive/>)<sup>25</sup>, via the ProteomeXchange initiative (<https://www.proteomexchange.org>)<sup>26</sup>.

Initiatives for establishing standards and recommendations of best practices within structural MS techniques, ion mobility-MS ([https://chemrxiv.org/articles/Recommendations\\_for\\_Reporting\\_Ion\\_Mobility\\_Mass\\_Spectrometry\\_Measurements/7072070](https://chemrxiv.org/articles/Recommendations_for_Reporting_Ion_Mobility_Mass_Spectrometry_Measurements/7072070)), hydrogen/deuterium exchange (manuscripts in preparation), and native MS are or have recently emerged. Likewise, there is also a clear need for the objective assessment of the methods and reporting standards within the field of XL-MS. For this purpose, several researchers active in the field of XL-MS decided to start a community-organized effort with the goal of providing a first overview of common procedures in XL-MS to generate the basis for best practices in the field.


In this first inter-laboratory effort, 32 groups worldwide contributed, delivering a total of 58 cross-linking data sets. The data reflect the great diversity of experimental and computational strategies employed and to our knowledge, this is the first comprehensive study with the aim to harmonize the XL-MS field.

## RESULTS.

*Study Design.* We opted for a simple study design to encourage participation from as many laboratories as possible, including those with currently only little experience in XL-MS. Invitations were sent out to research groups known to be active in the field from their published work and to attendants of the Symposium of Structural Proteomics (SSP, <http://www.structuralproteomics.net/>)

meeting series. The guidelines were kept quite simple and each participant was provided with a template spreadsheet to document their method and report their results (Supporting Information). Bovine serum albumin (BSA), a protein with a molecular weight of ~66 kDa, was selected as study system. We requested that a certain product from a widely available supplier should be used, and it was specified to use a BSA concentration of 10  $\mu$ M. Apart from these restrictions, we left the contributing labs full freedom to choose the experimental and computational strategies of their choice. This included, among other parameters, flexibility regarding the choice of cross-linking reagent and its concentration, buffer composition and pH, reaction time and temperature, post-cross-linking sample processing (digestion protocol, optional fractionation and enrichment of cross-linked products), conditions for LC/MS analysis, and data analysis procedures (choice of software, search parameters, validation of the results). In short, we expected that participants would use the typical XL-MS workflows established in their labs. The protocols used by the individual participating labs were collected and analyzed in the Sinz lab and are summarized in the Supporting Information.

For data analysis, we provided the amino acid sequence of mature BSA after cleavage of the signal peptide and propeptide sequences (residues 25-607 of the UniProt entry P02769; <https://www.uniprot.org/uniprot/P02769>) to ensure a uniform numbering scheme. Finally, we encouraged participants to perform at least three replicates. As mentioned above, we provided a template spreadsheet (Supporting Information) that needed to be completed by the participants before a data set would be considered for inclusion in the detailed assessment of the results. An overview of the data sets provided by different labs is presented in Figure 1.




**Figure 1.** Overview of data sets provided by the participants of this study. 32 groups participated in this study, yielding 58 separate cross-linking workflows. Nine datasets had to be excluded due to missing replicates and non-uniform conditions, resulting in a total of 49 datasets that were further considered. Several workflows contain both in-solution (47 samples) as well as in-gel digestion (10 samples) as processing methods. The samples were considered only once during a workflow analysis.

**Protein System.** BSA was selected as model protein for this study as it is a globular and stable protein that is readily available at low cost. Moreover, the three-dimensional structure of BSA is well-known and we selected the Protein Data Bank entry 4F5S

(<https://www.rcsb.org/structure/4F5S>) for further interpretation of the results. As BSA possesses a tendency towards forming dimers, this has to be considered when interpreting the results (see also below).

**Cross-linking Reagents.** As outlined above, the participants of this study were free to choose the cross-linking principle(s) on their own (Table S1; Supporting Information). The majority of groups decided to use non-cleavable, homobifunctional, amine-reactive N-hydroxysuccinimide (NHS) cross-linkers, i.e., bis(sulfosuccinimidyl)suberate (BS3) or disuccinimidylsuberate (DSS) (Figure 2a). Both cross-linkers only differ by a sulfonic acid group that is incorporated for increased water solubility and bridge a distance of 11.4 Å, resulting in C $\alpha$ -C $\alpha$  distances of ~27 Å to be cross-linked<sup>27</sup>. MS-cleavable cross-linkers, such as disuccinimidylsulfoxide (DSSO) and disuccinimidyl dibutyric urea (DSBU), are increasingly being used as they allow a targeted identification of cross-linked product based on characteristic reporter ions generated during MS/MS experiments. MS-cleavability as a cross-linker feature is essential to reduce the search space in conducting proteome-wide cross-linking studies. The vast majority of cross-linkers used herein target amine groups in proteins, i.e., lysine side chains, while carboxylic acid groups, such as aspartic and glutamic acid residues, are less frequently targeted (Figure 2b). The main spacer lengths of the cross-linkers are determined by the three most abundant cross-linkers used in this study are: BS3 and DSS (both 11.4 Å), DSBU (12.5 Å), and DSSO (10.1 Å) (Figure 2c).




**Figure 2.** (a) Cross-linking reagents used in this study; non-cleavable cross-linkers are presented in red, MS-cleavable cross-linkers are shown in blue; (b) reactivity, (c) spacer length. The cross-linkers used in this study are: BS<sup>3</sup> (bis(sulfosuccinimidyl)suberate), DSS (disuccinimidylsuberate), DSP (dithiobis(succinimidylpropionate)), DMTMM (4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methyl-morpholinium chloride) with and without PDH (pimelic acid dihydrazide), sulfo-SDA (sulfosuccinimidyl 4,4'-azipentanoate), CBSS (carboxybenzophenone sulfosuccinimide), DSSO (disuccinimidylsulfoxide), DSBU (disuccinimidyl dibutyric urea), BDP-NHP (N-hydroxyphthalimide ester of biotin aspartate proline), CBDPS (cyanurbiotindimercaptopropionyl succinimide), DC4 (1,4-bis(4-((2,5-dioxopyrrolidin-1-yl)oxy)-4-oxobutyl)-1,4-diazabicyclo[2.2.2]octane-1,4-diium), and MC4 (N,N'-bis(4-((2,5-dioxopyrrolidin-1-yl)oxy)-4-oxobutyl)-morpholine).

**Reaction Conditions.** The reaction conditions were also kept completely open to the participants, including cross-linking reaction time, temperature, cross-linker excess, and pH value of the cross-linking solution (Figure 3). Not surprisingly, the pH value of the


cross-linking reaction mixture was kept around pH 7.4 to 7.5 in the majority of experiments in order to resemble the physiological pH situation. A pH value of 8.0 that was also used in some experiments has the advantage of enhancing the reactivity of NHS esters with nucleophiles. The temperature was kept to 20, 25 or 37 °C in the majority of experiments, with lower temperature being applied only by a few groups. For BSA, a temperature of 37 °C certainly does not present a problem as it is a stable, globular protein, but for delicate and unstable proteins one should take care to conduct the cross-linking reaction at lower temperatures.

#### Instrument Platforms and Settings Used to Generate XL-MS Data.

The overwhelming majority of cross-linking data were generated on Orbitrap mass spectrometers (Figure 4). Only two FTICR (Solarix and Velos FTICR) mass spectrometers and one Q-TOF (Synapt G2 SI) instrument were employed (Figure 4a). All groups used LC/ESI-MS/MS analysis, applying for most experiments a resolving power of 60,000 or 120,000 (at  $m/z$  200 or 400, as specified by the manufacturer Thermo Fisher Scientific for orbitrap instruments) (Figure 4b). For MS/MS experiments, a resolving power of 15,000 or 30,000 was employed in most cases (Figure 4c). Details on enrichment of cross-linked species, considered charge states, fragmentation methods, and MS3 resolution are presented in the Supporting Information (Figure S1).




**Figure 3.** (a) Time, temperature, and cross-linker excess (XL-fold) were set as variable parameters, presented as grey spheres. The colored dots are projections of the 3D space onto 2D planes. (b) The pH values of the cross-linking reactions ranged between 7.0 to 8.2.



**Figure 4.** LC/MS/MS conditions applied. (a) MS instrumentation, (b) MS resolving power, (c) MS/MS resolving power. Resolving power is defined at  $m/z$  200 for orbitrap instruments, while for ICR instruments it is defined at  $m/z$  400. Please note that several research groups generated datasets with different instruments and settings.

**Data Analysis and Validation Strategies.** Strategies for data analysis were highly diverse (Figure 5), reflecting the variety in the XL-MS field where nearly every group possesses their own software tools tailored to fit their specific needs. This enormous variety is currently one of the most critical issues in XL-MS and we consider it as important contribution of this study to reflect this diversity. The false discovery rate (FDR) plays an important role in this context and from this study it arose that most of the groups apply an FDR of 5% (Figure 5b). Manual validation of the cross-links was performed for 66% of the experiments, while in 34%, the data sets were not manually checked. It is important to note that a mechanism to control the FDR should exist in the software, although proper FDR control is not trivial for small search spaces, manual validation strategies might be especially beneficial in such cases. Some strategies provide additional layers of evidence that can be used to better control the error rate. For example, isotope-coded, non-cleavable linkers provide two independent measures of precursor and fragment masses, and charge state information for fragments independent of MS resolution; MS-cleavable linkers provide three layers of information - intact precursors, released fragments corresponding to intact peptide chains, and fragments thereof. In absence of such strategies, we recommend that preferentially both, MS and MS/MS data, should be recorded with high mass accuracy to rule out a false assignment of cross-linked products. Clearly, some of these effects will only become apparent for samples of higher complexity.






**Figure 5.** (a) Software tools used in this study (a complete summary is found in Table S2; Supporting Information). Red bars indicate that the software is applicable only for non-cleavable cross-linkers; blue bars indicate that the software can be used for MS-cleavable cross-linkers. (b) False discovery rates. (c) Mass tolerances MS. (d) Mass tolerances MS/MS. For the Proteome Discoverer, data analysis was performed using the XlinkX software node.

**Identified Cross-links.** As we left it to the individual participants whether to use in-solution or in-gel digestion as work-up method before LC/MS/MS analysis, 47 data sets were generated by in-solution digestion, while 10 samples originated from in-gel digestion (Figure 1). As already mentioned, BSA has a tendency to form dimers, which somewhat complicates data analysis. In case only the BSA monomer band is used for in-gel digestion and subsequent generation of the cross-linking data set, one can definitely rule out that cross-links are in fact representing intermolecular interactions between two BSA molecules. On the other hand, during the in-gel digestion procedure cross-links might get lost, resulting in an overall lower number of cross-linked products.

Another aspect regards the reaction sites that were considered during data analysis. Usually, NHS esters, such as the mainly used cross-linkers BS3, DSS, DSB, and DSSO will react with lysine, but they also exhibit a significant reactivity towards serine, threonine and tyrosine. The pH used for conducting the cross-linking reaction plays a significant role as amine reactivity is increased at higher pH values. Some participants considered only Lys-Lys cross-links and neglected the side-reactivity of NHS esters with hydroxy group-containing amino acids. In this study, it became apparent that Ser, Thr, and Tyr account for ~30% of cross-linking sites (Supporting Information, Figure S2). The reactivity of Ser, Thr, and Tyr residues obviously depends on the reaction conditions (cross-linker, pH value of the solution) as well as local pKa value. It is not practicable to consider Lys, Ser, Thr, and Tyr when analyzing very complex systems, such as complete proteomes. Therefore we suggest as a compromise to consider for whole proteome samples only lysine as reactive sites of NHS ester cross-linkers, while for single proteins or proteins assemblies, Lys, Ser, Thr, and Tyr might be taken into account.


Figure 6 provides an overview about the reproducibility of results obtained with the individual workflows of the participants. For in-solution digestion workflows, the average number of unique cross-links in BSA is 78, while for in-gel digestion workflows using only the monomeric BSA band, the average number is 44. The term “cross-link” refers to the specific amino acid residues that are connected, irrespective of different peptide sequences due to missed cleavage sites or modifications. The majority of participating labs

came up with similar numbers of unique cross-links, independently of the cross-linking conditions used (Figure 6a). Three cross-linking workflows however recorded a significantly higher number of cross-links (between 260 and 350). The reason could be a false consideration of cross-links from BSA dimers that in some preparations might have been a dominating species due to inappropriate sample treatment. For in-gel digestion workflows, up to 19 overlength cross-links were reported in one dataset, which could represent false-positives due to partial unfolding as only the monomeric form of BSA was considered in these samples (Figure 6b).



**Figure 6.** Number of BSA cross-links identified. The numbers of cross-links are plotted for (a) in-solution, (b) in-gel digestion workflows. The different cross-linkers are shown as symbols; abbreviations of the cross-linkers are according to Figure 2. The maximum distances are given for each cross-linker, indicating the number of overlength cross-links. Every point is a sum of three replicate measurements; replicates of the entire experiment are shown in blue, replicates of the LC/MS analyses are shown in red; the average number and reproducibility of unique cross-links are shown in yellow.

A more detailed inspection of the unique cross-links revealed highly interesting insights: Datasets created from amine-reactive cross-linkers (BS3, DSB, DSS, DSSO, DC4, MC4, CBDPS) using an in-solution digestion workflow yielded a total of 1066 unique cross-links. A complete list of unique cross-links, identified with cross-linkers reacting with nucleophiles (amine and hydroxy groups) and sorted by their reproducibility, is provided as separate file in the Supporting Information. 601 of 1066 unique cross links (56%) were however identified in only one single dataset (Figure 7). This indicates an overall low reproducibility of cross-linking results. The curve in Figure 7a shows that the number of unique cross-links identified is inversely proportional to the reproducibility of cross-links in the data sets (coefficient of proportionality  $\approx -1$ ). If the reproducibility across the data sets is higher than 20%, the effect of including more datasets, different reaction conditions, and analytical parameters determines a linear increment of the number of cross-link identifications. The intercept with the y-axis of the resulting interpolated linear curves indicates the putative number of cross-links in BSA to be between 73 and 88 (Figures 7b). This value is very close to the average number of cross-links found (78 cross-links per dataset for in-solution digestion workflows, Figure 6a). In Figure 7c, the dependence of the linear correlation on the reproducibility of cross-links identified is indicated. This indicates that a linear correlation only exists for highly reproducible cross-links.



**Figure 7.** Comparison of unique cross-links. “Cross-link” denotes the unique amino acid residues that are connected by homobifunctional, amine-reactive cross-linkers. (a) Number of cross-links in respect to their reproducibility among the data sets. (b) Linear extrapolation of all (red) or a linear subset (blue) of cross-links resulted in a maximum cross-linking number between 73 and 88. (c) The plot summarizes the intercepts with the y-axis (red) and the correlation coefficient  $\times 100$  (blue) of the respective linear extrapolations of (a). The linear extrapolation was calculated as shown in (b) by successively removing the data points starting from the lowest reproducible value.

*Cross-links identified from in-gel digested BSA monomer band.* We mapped cross-links in the monomer band of BSA using *in-gel* digestion (in total 10 datasets) into the published 3D structure of BSA (pdb entry 4F5S). For this, homobifunctional, amine-reactive linkers were considered as only for this type of cross-linkers a statistical analysis could be performed. Only cross-links identified in at least two independent experiments are presented (Figure S3). 30 out of 230 cross-links exceed the maximum length of 30 Å for the cross-linkers employed in this study. These overlength cross-links either originate from a false assignment or either by applying non-suitable experimental conditions. Strikingly, 29 of these overlength cross-links were identified in one single experiment only. Cross-links that were identified in at least two independent experiments show one over-length link, while cross-links found in at least three independent experiments all fall within the given distance limit of 30 Å (Figure S4). As guideline for testing cross-linking workflows, we provide a list of cross-links that were identified in at least two independent experiments from *in-gel* digestion of the BSA monomer band (Table S3; Supporting Information).

*Monomer-dimer equilibrium of BSA.* BSA exists in a monomer-dimer equilibrium, which may give rise to ambiguities in the identification of intra- and inter-molecular cross-link. To address this issue, we performed additional experiments with four concentrations of BSA (10, 5, 1, and 0.5  $\mu\text{M}$ ). Strikingly, the number of overlength cross-links was very low (only 1 or 2). Moreover, the numbers of overlength cross-links were similar for all four BSA concentrations used (Table S4; Supporting Information). This clearly indicates that a BSA concentration of 10  $\mu\text{M}$ , as chosen for this study is suitable for conducting cross-linking MS experiments.

*Comparison of data acquisition and analysis strategies from one participating lab.* Because most of the data in this study have been gener-

ated in different laboratories, differences in instrumentation and in the software used for data analysis make a direct comparison of selected results difficult. However, we used a subset of the data generated in a single laboratory to study the effect of the type of mass spectrometer and of different search settings on the outcome for a relatively simple model system, such as BSA (see Supporting Information).

## DISCUSSION.

This first community-based cross-linking study reflects the high diversity of XL-MS workflows that are currently employed in different labs worldwide. However, it also became apparent that independently of the workflow used the results obtained are to some degree comparable. For beginners in the field we suggest to use BSA as an initial study system and compare the outcome to the results obtained herein. As a guideline, the number of cross-links expected for BSA should be  $\sim 80$  for an *in-solution* workflow, considering cross-links of the monomer and the dimer. Not unexpectedly, our study did not reveal the optimum experimental protocol or software to be used in any and all projects. The applications of XL-MS are just too diverse so that no single cross-linker, instrument or software tool is expected to be preferable for all scenarios, ranging from single protein (as used in this work) to whole-cell cross-linking. There are also clear interdependencies between the type of cross-linker (cleavable, non-cleavable) and the software that can be applied to process such data, as well as between instrument type and software as not all fragmentation methods or other MS platform-dependent features may be supported.

As discussed above, XL-MS has become an essential part of many structural proteomics studies, but is also a key element in integrative structural biology projects. In such interdisciplinary work, XL data may only be a small “puzzle piece” that is combined with other experimental data provided by methods such as electron microscopy, X-ray crystallography, NMR spectroscopy, small-angle X-ray scattering, together with computational modeling. Details about how experiments were carried out, how the data were processed, and how error rates were assessed are often missing from the publication, making it difficult for reviewers and readers to assess the reliability and credibility of the results. We therefore recommend that appropriate consideration should be given to the method section of all XL-MS publications by providing all necessary experimental and computational details. Our reporting template could serve as a starting point for the “minimum information about a cross-linking experiment” that should be included in research articles containing XL-MS data. This template is included in the Supporting Information for all XL-MS data reports. Sufficient information needs to be provided, irrespective of the relative contribution of the cross-linking experiments to a specific project. This will also facilitate the cross-referencing of XL-MS data in integrative structural biology projects, for example in the dedicated PDB prototype archive, PDB-Dev<sup>28</sup>.

Data deposition to a proteomics repository, such as PRIDE, is encouraged, as the paucity of available data sets do not assist the field in validation, methods evaluation and workflow quality. It should be noted that not all data sets assigned to the cross-linking category in PRIDE originate from genuine XL-MS experiments (in the sense that cross-linking sites were identified), but also contain



data from experiments that used cross-linking for the stabilization of complexes. The low uptake of data deposition may in part be due to the specific nature of XL-MS data. For a “complete” submission to ProteomeXchange, allowing a complete integration of search results and assignment of a Digital Object Identifier, the reported results need to be compliant with a PSI format, such as mzIdentML. Although the most recent version of mzIdentML

In summary, seven guidelines were deduced from this study as framework for conducting XL-MS experiments as detailed in Table 1.

Table 1. Cross-linking mass spectrometry guidelines. Guidelines 1 and 2 are derived from the results shown in Figure 5.

#	Topic	Description
1	FDR	A mechanism to control the FDR should exist in the software used for cross-link identification. The FDR algorithm has to be described in detail. For small search spaces, manual validation strategies might be beneficial.
2	Mass accuracy	MS and MS/MS data should be recorded and analyzed with high mass accuracy to reduce false assignments of cross-linked products, or multiple lines of evidence from isotope labeling or cleavable linkers should be obtained.
3	Experimental details	Provide all experimental and computational details. The reporting template (Supporting Information) comprises the “minimum information of a cross-linking experiment” that should be included in research articles containing XL-MS data.
4	Data deposition	Deposit raw MS files together with a description of their content and the reporting template to a proteomics repository, such as PRIDE.
5	Visualization of cross-linked proteins	Perform SDS-PAGE analysis to evaluate the cross-linking performance under the employed experimental conditions. Check for possible high-molecular weight aggregates.
6	Cross-linker selectivity	Consider only lysine and the N-terminus as reactive sites of amine-reactive cross-linkers for whole proteome samples. For single proteins or large protein assemblies, consider lysine, N-terminus, serine, threonine, and tyrosine as reactive sites.
7	BSA cross-links	Approximately 80 cross-links can be expected for cross-linking of BSA using homobifunctional amine-reactive cross-linkers and an in-solution digestion workflow.

(version 1.2) includes support for some XL-MS strategies, such as proteomics-centered format cannot easily consider all possible workflows, and few dedicated cross-linking search engines offer mzIdentML-compliant export at this point. Nevertheless, even a “partial” submission will make the raw MS data and results available in a user-specified format for download and re-use by interested researchers.

Additional studies that cover a wider range of sample types, such as large multi-protein assemblies or even whole proteomes, will be required to obtain a better understanding of the benefits and drawbacks of different experimental workflows. However, we believe that this first community-based study serves as the starting point for further initiatives in this direction, and encourages the adoption of consistent reporting and data sharing guidelines in XL-MS. We would like to invite interested parties to participate in the discussion to expand the growing XL-MS community.

## CONCLUSION AND GUIDELINES

Although XL-MS is becoming routine for in vitro and in vivo applications in proteomics and structural biology, this harmonization initiative unveiled a great variety in the cross-links identified by participating groups, even for the single protein BSA. This underlines the need for establishing generally accepted XL-MS protocols as well as data analysis and reporting formats. This inter-laboratory study on XL-MS represents the first effort of the community towards establishing endorsed and transparent good practice guidelines for performing and reporting XL-MS experiments. This study also serves as test for all laboratories to evaluate the quality of their XL-MS workflows and will aid in improving eventual weaknesses.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS website. The workflows employed by the participating groups are presented in a spreadsheet. The cross-links identified with homobifunctional, amine-reactive cross-linkers (*in-solution* digestion) are presented. The cross-links identified with homobifunctional, amine-reactive cross-linkers (*in-gel* digestion, monomer band) in at least two independent experiments are presented.

## AUTHOR INFORMATION

### Corresponding Authors

\* Andrea Sinz, Department of Pharmaceutical Chemistry & Bioanalytics, Institute of Pharmacy, Charles Tanford Protein Center, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Str. 3a, D-06120 Halle (Saale), Germany.

Email address: [andrea.sinz@pharmazie.uni-halle.de](mailto:andrea.sinz@pharmazie.uni-halle.de)

\* Alexander Leitner, Institute of Molecular Systems Biology, Otto-Stern-Weg 3, ETH Zurich, CH-8093 Zurich, Switzerland.

Email address: [leitner@imsb.biol.ethz.ch](mailto:leitner@imsb.biol.ethz.ch)

# Both authors contributed equally.

### Author Contributions

The study was designed by AL, CI, and AS, all authors contributed data. CI, CP, AL, and AS performed data analysis. SF coordinated the COST Action BM1403.

## ACKNOWLEDGMENT

## REFERENCES

- (1) Young, M. M.; Tang, N.; Hempel, J. C.; Oshiro, C. M.; Taylor, E. W.; Kuntz, I. D.; Gibson, B. W.; Dollinger, G. High throughput protein fold identification by using experimental constraints derived from intramolecular cross-links and mass spectrometry. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 5802-5806 (2000).
- (2) Rappsilber, J.; Siniosoglou, S.; Hurt, E. C.; Mann, M. A Generic Strategy To Analyze the Spatial Organization of Multi-Protein Complexes by Cross-Linking and Mass Spectrometry. *Anal. Chem.* **2000**, *72*, 267-275.
- (3) Bennett, K. L.; Kussmann, M.; Björk, P.; Godzwon, M.; Mikkelsen, M.; Sørensen, P.; Roepstorff, P. Chemical cross-linking with thiol-cleavable reagents combined with differential mass spectrometric peptide mapping--a novel approach to assess intermolecular protein contacts. *Protein Sci.* **2000**, *9*, 1503-1518.
- (4) Sinz, A.; Wang, K. Mapping Protein Interfaces with a Fluorogenic Cross-Linker and Mass Spectrometry: Application to Nebulin-Calmodulin Complexes. *Biochemistry* **2001**, *40*, 7903-7913.
- (5) Tang, X.; Munske, G. R.; Siems, W. F.; Bruce, J. E. Mass spectrometry identifiable cross-linking strategy for studying protein-protein interactions. *Anal. Chem.* **2005**, *77*, 311-318.
- (6) Leitner, A.; Walzthoeni, T.; Aebersold, R. Lysine-specific chemical cross-linking of protein complexes and identification of cross-linking sites using LC-MS/MS and the xQuest/xProphet software pipeline. *Nat. Protoc.* **2014**, *9*, 120.
- (7) Schmidt, C.; Robinson, C. V. A comparative cross-linking strategy to probe conformational changes in protein complexes. *Nat. Protoc.* **2014**, *9*, 2224.
- (8) Lima, D. B.; Melchior, J. T.; Morris, J.; Barbosa, V. C.; Chamot-Rooke, J.; Fioramonte, M.; Souza, T. A. C. B.; Fischer, J. S. G.; Gozzo, F. C.; Carvalho, P. C.; Davidson, W. S. Characterization of homodimer interfaces with cross-linking mass spectrometry and isotopically labeled proteins. *Nature protocols* **2018** *13*, 431-458.
- (9) Orbán-Németh, Z.; Beveridge, R.; Hollenstein, D. M.; Rampler, E.; Stranzl, T.; Hudecz, O.; Doblmann, J.; Schlögelhofer, P.; Mechtler, K. Structural prediction of protein models using distance restraints derived from cross-linking mass spectrometry data. *Nat. Protoc.* **2018**, *13*, 478-494.
- (10) Liu, F.; Lössl, P.; Scheltema, R.; Viner, R.; Heck, A. Optimized fragmentation schemes and data analysis strategies for proteome-wide cross-link identification. *J. Nat. Commun.* **2017**, *8*, 15473.
- (11) Klykov, O.; Steigenberger, B.; Pektaş, S.; Fasci, D.; Heck, A. J.; Scheltema, R. A. Efficient and robust proteome-wide approaches for cross-linking mass spectrometry. *Nat. Protoc.* **2018**, *13*, 2964-2990.
- (12) Iacobucci, C.; Götz, M.; Ihling, C. H.; Piotrowski, C.; Arlt, C.; Schäfer, M.; Hage, C.; Schmidt, R.; Sinz, A. A cross-linking/mass spectrometry workflow based on MS-cleavable cross-linkers and the MeroX software for studying protein structures and protein-protein interactions. *Nat. Protoc.* **2018**, *13*, 2864-2889.
- (13) Chen, Z. A.; Rappsilber, J. Quantitative cross-linking/mass spectrometry to elucidate structural changes in proteins and their complexes. *Nat. Protoc.* **2018**, *14*, 171-201.
- (14) Bell, A. W.; Deutsch, E. W.; Au, C. E.; Kearney, R. E.; Beavis, R.; Sechi, S.; Nilsson, T.; Bergeron, J. J.; HUPO Test Sample Working Group. A HUPO test sample study reveals common problems in mass spectrometry-based proteomics. *Nat. Methods* **2009**, *6*, 423.
- (15) Addona, T. A.; Abbatiello, S. E.; Schilling, B.; Skates, S. J.; Mani, D. R.; Bunk, D. M.; Spiegelman, C. H.; Zimmerman, L. J.; Ham, A. J.; Keshishian, H.; et al. Multi-site assessment of the precision and reproducibility of multiple reaction monitoring-based measurements of proteins in plasma. *Nat. Biotech.* **2009**, *27*, 633.
- (16) Abbatiello, S. E.; Schilling, B.; Mani, D. R.; Zimmerman, L. J.; Hall, S. C.; MacLean, B.; Albertolle, M.; Allen, S.; Burgess, M.; Cusack, M. P.; et al. Large-Scale Interlaboratory Study to Develop, Analytically Validate and Apply Highly Multiplexed, Quantitative Peptide Assays to Measure Cancer-Relevant Proteins in Plasma. *Mol. Cell. Proteomics* **2015**, *14*, 2357-2374.
- (17) Kennedy, J. J.; Abbatiello, S. E.; Kim, K.; Yan, P.; Whiteaker, J. R.; Lin, C.; Kim, J. S.; Zhang, Y.; Wang, X.; Ivey, R. G.; et al. Demonstrating the feasibility of large-scale development of standardized assays to quantify human proteins. *Nat. Methods* **2013**, *11*, 149.
- (18) Vialas, V.; Colomé-Calls, N.; Abian, J.; Aloria, K.; Alvarez-Llamas, G.; Antúnez, O.; Arizmendi, J. M.; Azkargorta, M.; Barceló-Batllo, S.; Barderas, M. G.; et al. A multicentric study to evaluate the use of relative retention times in targeted proteomics. *J. Proteomics* **2017**, *152*, 138-149.
- (19) Navarro, P.; Kuharev, J.; Gillet, L. C.; Bernhardt, O. M.; MacLean, B.; Röst, H. L.; Tate, S. A.; Tsou, C. C.; Reiter, L.; Distler, U.; et al. A multicenter study benchmarks software tools for label-free proteome quantification. *Nat. Biotech.* **2016**, *34*, 1130 (2016).
- (20) Collins, B. C.; Hunter, C. L.; Liu, Y.; Schilling, B.; Rosenberger, G.; Bader, S. L.; Chan, D. W.; Gibson, B. W.; Gingras, A. C.; Held, J. M.; et al. Multi-laboratory assessment of reproducibility, qualitative and quantitative performance of SWATH-mass spectrometry. *Nat. Commun.* **2017**, *8*, 291.
- (21) Deutsch, E. W.; Orchard, S.; Binz, P. A.; Bittremieux, W.; Eisenacher, M.; Hermjakob, H.; Kawano, S.; Lam, H.; Mayer, G.; Menschaert, G.; et al. Proteomics Standards Initiative: Fifteen Years of Progress and Future Work. *J. Proteome Res.* **2017**, *16*, 4288-4298.
- (22) Taylor, C. F.; Paton, N. W.; Lilley, K. S.; Binz, P. A.; Julian, R. K. Jr.; Jones, A. R.; Zhu, W.; Apweiler, R.; Aebersold, R.; Deutsch, E. W.; et al. The minimum information about a proteomics experiment (MIAPE). *Nat. Biotech.* **2007**, *25*, 887.
- (23) Martens, L.; Chambers, M.; Sturm, M.; Kessner, D.; Levander, F.; Shofstahl, J.; Tang, W. H.; Römpp, A.; Neumann, S.; Pizarro, A. D.; et al. mzML--a community standard for mass spectrometry data. *Mol. Cell. Proteom.* **2011**, *10*, R110.000133.
- (24) Vizcaino, J. A.; Mayer, G.; Perkins, S.; Barsnes, H.; Vaudel, M.; Perez-Riverol, Y.; Ternent, T.; Uszkoreit, J.; Eisenacher, M.; Fischer, L.; et al. The mzIdentML Data Standard Version 1.2, Supporting Advances in Proteome Informatics. *Mol. Cell. Proteom.* **2017**, *16*, 1275-1285 (2017).
- (25) Vizcaino, J. A.; Csordas, A.; del-Toro, N.; Dianes, J. A.; Griss, J.; Lavidas, I.; Mayer, G.; Perez-Riverol, Y.; Reisinger, F.; Ternent, T.; et al. 2016 update of the PRIDE database and its related tools. *Nuc. Acids Res.* **2015**, *44*, D447-D456.
- (26) Deutsch, E. W.; Csordas, A.; Sun, Z.; Jarnuczak, A.; Perez-Riverol, A.; Ternent, T.; Campbell, D. S.; Bernal-Llinares, M.; Okuda, S.; Kawano, S.; et al. The ProteomeXchange consortium in 2017: supporting the cultural change in proteomics public data deposition. *Nuc. Acids Res.* **2016**, *45*, D1100-D1106.
- (27) Merkley, E. D.; Rysavy, S.; Kahraman, A.; Hafen, R. P.; Daggett, V.; Adkins, J. N. Distance restraints from crosslinking mass spectrometry: mining a molecular dynamics simulation database to evaluate lysine-lysine distances. *Protein Sci.* **2014**, *23*, 747-759.
- (28) Vallat, B.; Webb, B.; Westbrook, J. D.; Sali, A.; Berman, H. M. Development of a Prototype System for Archiving Integrative/Hybrid Structure Models of Biological Macromolecules. *Structure* **2018**, *26*, 894-904.

## 1<sup>ST</sup> INTERLABORATORY XLMS STUDY

32 groups participating worldwide

